arXiv:quant-ph/0604047v1 7 Apr 2006

Characterizing quantum phase transitions by single qubit operations

S. M. Giampaolo, F. Illuminati, and S. De Siena
Dipartimento di Fisica "E. R. Caianiello", Università di Salerno, Coherentia CNR-INFM,
and INFN Sezione di Napoli - Gruppo Collegato di Salerno, Via S. Allende, 84081 Baronissi (SA), Italy
(Dated: April 7, 2006)

We introduce observable quantities, borrowing from concepts of quantum information theory, for the characterization of quantum phase transitions in spin systems. These observables are uniquely defined in terms of single spin unitary operations. We define the energy gap between the ground state and the state produced by the action of a single-qubit local gate. We show that this static quantity involves only single-site expectations and two-point correlation functions on the ground state. We then discuss a dynamical local observable defined as the acceleration of quantum state evolution after performing an instaneous single-qubit perturbation on the ground state. This quantity involves three-point correlations as well. We show that both the static and the dynamical observables detect and characterize completely quantum critical points in a class of spin systems.

PACS numbers: 03.67.-a, 75.10.Jm, 05.50.+q

Quantum phase transitions differ from classical ones in that they are driven by quantum fluctuations. They are induced by the changes in external control parameters or coupling constants, which can modify the ground state of a quantum system [1]. In analogy with classical phase transitions driven by thermal fluctuations, quantum phase transitions are usually analyzed in terms of the scaling behavior of equilibrium properties and diverging correlation lengths as signatures of quantum critical points [1, 2]. In the last years interest in quantum critical phenomena has increased enormously thanks to several studies that have highlighted some intriguing connections between quantum phase transitions and the scaling behavior of nonlocal quantum correlations in the ground state of complex quantum systems [3, 4, 5]. Quantum critical points in some classes of quantum spin models can be characterized by the divergence in the first derivative of the two-site (two-qubit) concurrence, a proper measure of bipartite entanglement [6], with respect to the proper control parameter, even though recent work on different systems has revealed that this behavior of the concurrence is not universal at generic quantum critical points [7, 8, 9], and different classification schemes have been proposed recently [10, 11]. The studies on the scaling behavior of entanglement prompt naturally the question on the possible links of entanglement measures to observable quantities in quantum critical systems. More generally, as quantum entanglement plays a fundamental role both in quantum information and quantum computation [12], the further question arises whether it is possible to reveal structural aspects of quantum critical systems using other concepts and techniques, beyond entanglement, motivated and/or imported from the arena of quantum information theory.

Following this line of thought, in this work we discuss the behavior of observables associated to unitary transformations on individual spins, i.e. local unitary single qubit operations. We will show that they can be used to detect and characterize quantum critical points in a class of quantum spin systems. First, we will focus our attention on the energy gap needed to realize single qubit operations by local excitations above the ground state. Next, we will consider the short-time diffusion coefficient of the on site magnetization along a given direction, i.e. the quantum speed characterizing the dynamics of the magnetization under the action of single qubit operations. We will first establish that the observable quantities associated to fundamental local operations (elementary local gates) can be unambiguously expressed as functions of one-, two-, and three-point correlation functions evaluated in the ground state. These relations in turn will allow to study their behavior when approaching a quantum phase transition, and to detect and characterize it.

Let us consider the class of spin-1/2 ferromagnetic systems described by the Hamiltonian

$$H=-\frac{J}{2}\sum_i\left[(1+\gamma)\sigma_i^x\sigma_{i+1}^x+(1-\gamma)\sigma_i^y\sigma_{i+1}^y\right]+h\sum_i\sigma_i^z\,, \tag{1}$$

where i runs over the sites of the lattice, J is the nearest-neighbor coupling constant, h is the transverse magnetic field directed along the z-axis, γ is the anisotropy parameter, $0\le\gamma\le1$, σ_i^α are the Pauli matrices ($\alpha=x,\,y,\,z$), and periodic boundary conditions are assumed throughout. It is convenient to define the dimensionless reduced coupling constant $\lambda=J/h$. These models are exactly solvable [13, 14, 15], and, in principle, the structure of the correlation functions in the ground state can be studied accurately. For $\gamma=1$ Eq. (1) reduces to the Ising model whereas for $\gamma=0$ it reduces to the isotropic XY model. In the interval $0<\gamma\le1$ the models belong to the Ising universality class. For $0<\gamma\le1$, in the thermodynamic limit the system undergoes an order-disorder phase transition at the critical value $\lambda_c=1$. If $\lambda>\lambda_c$ the magnetization $<\sigma_i^x>$ is finite and vanishes at the transition, while the magnetization $<\sigma_i^z>$ remains finite for any value of λ and vanishes only in the limit $\lambda\to\infty$.

The case $\gamma = 0$ is in a critical regime for all $\lambda \geq \lambda_c$.

A local gate (or one qubit gate) is any unitary transformation acting as the identity $\mathbf{1}$ on any subspace defined on any spin of the lattice except one. Consider the single spin, on which the local gates act nontrivially, located, say, at a certain site k. Then, the corresponding unitary operator can be written as $U = \prod_{i \neq k} \mathbf{1}_i O_k$, where O_k is a generic unitary operation acting on the spin placed on site k [12]. The natural basis in the space of local gates at lattice site k is formed by the three Pauli matrices σ_k^x, σ_k^y, σ_k^z, and the identity $\mathbf{1}_k$. For instance, the Hadamard gate is $H_k = (\sigma_k^x + \sigma_k^z)/\sqrt{2}$, the phase gate is $S_k = ((1-i)\sigma_k^z + (1+i)\mathbf{1}_k)/2$, and the spin flip operator is simply σ_k^x.

The action of a local gate on the ground state $|g\rangle$ of the system takes it to a state $|\psi\rangle = O_k|g\rangle$ that, in general, is not an eigenstate of the many-body Hamiltonian H. Introducing the (dimensionless) energy gap $\Delta E(O_k) = (\langle\psi|H|\psi\rangle - \langle g|H|g\rangle)/h$ it is possible to determine the exact expressions for the three energy gaps associated to the fundamental gates in terms of various correlation functions. We obtain:

$$\begin{aligned} \Delta E_x &= 2\lambda(1-\gamma)G_{yy} - 2M_z \ , \\ \Delta E_y &= 2\lambda(1+\gamma)G_{xx} - 2M_z \ , \\ \Delta E_z &= 2\lambda\left[(1+\gamma)G_{xx} + (1-\gamma)G_{yy}\right] \ , \end{aligned} \quad (2)$$

where ΔE_α denotes the energy gap due to the action of the gate σ_k^α, M_z is the value of the on site magnetization along the z-axis in the ground state: $M_z = \langle g|\sigma_k^z|g\rangle$, and $G_{\alpha\alpha}$ denote the values of the nearest neighbor two-point correlation functions in the ground state: $G_{\alpha\alpha} = \langle g|\sigma_k^\alpha \sigma_{(k+1)}^\alpha|g\rangle$. In the derivation of Eqs. (2) we properly exploited the invariance of Hamiltonian Eq. (1) under translations.

The major significance of the set of Eqs. (2) is that the energy gaps between excited states obtained by local gate operations on the ground state and the ground state itself can be expressed as functions of (bipartite) expectation values on $|g\rangle$. Therefore, correlation functions and other ground state properties can be determined once the energy gaps entering in Eqs. (2) are known, and changes in $|g\rangle$ must be reflected in changes of the energy gaps.

In Fig.1 we plot the behavior of the derivatives with respect to λ of the three energy gaps in the case of the Ising model ($\gamma = 1$). Here and in the following of this work we will exploit the studies performed by Barouch and McCoy [14] to obtain exact analytical expressions. We find that the derivatives of the three energy gap diverge logarithmically at the critical point $\lambda_c = 1$ as

$$\frac{d\Delta E_\alpha}{d\lambda} = c_\alpha(\gamma) \ln|\lambda - \lambda_c| + const \ , \quad (3)$$

where, for the Ising model, the values of the coefficients $c_\alpha(1)$ are $c_z(1) = -1.276$, $c_x(1) = 0.638$, and $c_y(1) = -c_x(1)$.



FIG. 1: First derivatives of the energy gaps as functions of λ in the Ising model ($\gamma = 1$). Red line is $d\Delta E_z/d\lambda$, blue line $d\Delta E_y/d\lambda$, and black line $-d\Delta E_x/d\lambda$.

For $0 < \gamma < 1$ the models described in Eq. (1) are in the same universality class of the Ising model. With respect to the concept of universality - the critical properties depend only on the dimensionality of the system and on the broken symmetry in the ordered phase - it is important to verify that the energy gaps show similar behaviors when the models in this range of values of γ approach the quantum critical point. We have verified the same property of logarithmic divergence when the anysotropy parameter varies in the open range $(0, 1)$. For instance, for $\gamma = 0.5$ the derivatives of the energy gaps close to the quantum critical point obey the same law Eq. (3) with $c_z(0.5) = -2.550$, $c_x(0.5) = 0.638$ and $c_y(0.5) = -c_x(0.5)$. In general, for decreasing γ, we find that the value of c_z decreases and, remarkably, that the values of c_x and c_y, and the relation between them, remain constant.

Considering the isotropic XY model ($\gamma = 0$), the behavior of the energy gaps close to the quantum critical point changes radically because the system belongs to a universality class characterized by a new symmetry, the invariance under rotations around the z-axis. This symmetry imposes the plane degeneracy $E_x = E_y$ in Eqs. (2). In Fig.2 we show the behavior of the derivatives of the two remaining independent energy gaps (E_x and E_z) as a function of λ approaching the critical value λ_c. One finds that the derivative of the energy gap ΔE_z associated to the π phase gate operation σ_k^z diverges as an inverse power approaching λ_c from above:

$$\frac{d\Delta E_z}{d\lambda} = \frac{8}{\pi}(\lambda^2 - \lambda_c^2)^{-1/2} \ . \quad (4)$$

The derivative of the energy gap ΔE_x associated to the spin flip gate remains finite for all values of λ. Hence, only the energy gap associated to the (π) phase gate operation detects the onset of the critical regime in the XY model, due to the larger symmetry constraints.

Next, we consider quantum state dynamics rather than energy observables. In general, a local gate operation perturbs the ground state $|g\rangle$, and under the subsequent



FIG. 2: First derivatives of the two independent energy gaps as a function of λ for the XY model ($\gamma = 0$). Red line: $d\Delta E_z/d\lambda$; black line: $d\Delta E_x/d\lambda = d\Delta E_y/d\lambda$.



FIG. 3: First derivatives of the short-time diffusion coefficients Λ_x (black line), $-\Lambda_y$ (blue line), and Λ_z (red line), as functions of λ for the Ising model ($\gamma = 1$). Notice that Λ_z is always constant as a function of λ.

dynamics driven by Eq. (1) any observable that does not commute with the system Hamiltonian acquires a nontrivial time evolution. It is convenient to focus on the dynamics of the magnetization along the z-axis on site k, because it is nonvanishing for all values of λ, and hence its time evolution is best suited for analytical treatment as well as for experimental observation. Let us then define the dimensionless time $\tau = (ht)/\hbar$, and consider the short-time dynamics ($\lambda\tau \ll 1$) of the magnetization on site k after the local gate operation has been performed at $\tau = 0$. One has $< \sigma_k^z(\tau) > - < \sigma_k^z(0) >= -4\Lambda_\alpha(\lambda\tau)^2$, where $< \sigma_k^z(0) >$ is the magnetization along the z-axis evaluated on the perturbed state at time 0, $< \sigma_k^z(\tau) >$ is the on site magnetization at short times τ after the action of the local gate, and Λ_α is the short-time diffusion coefficient, or acceleration, whose form depends on the single qubit operation that has been performed. This acceleration can be expressed, similarly to the case of the energy gaps previously studied, as a linear combination of correlation functions evaluated in the ground state. For the three fundamental operations (σ_k^x, σ_k^y, and σ_k^z), the coefficients Λ_α can be worked out analytically and read, respectively,

$$\begin{aligned}
\Lambda_x &= -M_z + \frac{\gamma}{\lambda}(G_{xx} + G_{yy}) - \\
&\quad - \frac{1+\gamma}{2} G_{xzx} + \frac{\gamma(1-\gamma)}{2} G_{yzy} \; , \\
\Lambda_y &= -M_z - \frac{\gamma}{\lambda}(G_{xx} + G_{yy}) - \qquad (5) \\
&\quad - \frac{\gamma(1+\gamma)}{2} G_{xzx} - \frac{(1-\gamma)}{2} G_{yzy} \; , \\
\Lambda_z &= \gamma^2 M_z - \frac{\gamma}{\lambda}(G_{xx} - G_{yy}) \\
&\quad + \frac{1+\gamma}{2} G_{xzx} + \frac{1-\gamma}{2} G_{yzy} \; .
\end{aligned}$$

Comparing Eqs. (2) and Eqs. (5), we see that in the latter there appear three-point correlations involving nearest neighbor and next to nearest neighbor sites: $G_{\alpha\beta\alpha} = \langle g|\sigma_{k-1}^\alpha \sigma_k^\beta \sigma_{k+1}^\alpha|g\rangle$. The presence of correlations of order $n > 2$ is related to the dynamical nature of the observable. Including next-to-leading order powers of $\lambda\tau$ in the expansion of $< \sigma_k^z(\tau) > - < \sigma_k^z(0) >$, one would need to evaluate four-point and higher-order correlations.

In Fig.3 we plot the behavior of the derivatives of the diffusion coefficients $d\Lambda_\alpha/d\lambda$ ($\alpha = x, y, z$) as functions of λ for the Ising model ($\gamma = 1$). The derivatives of Λ_x and Λ_y diverge logarithmically approaching λ_c, in analogy with the derivatives of the energy gaps (see Eq. (3) and Fig.1), as

$$\frac{d\Lambda_\alpha}{d\lambda} = d_\alpha(\gamma) \ln|\lambda - \lambda_c| + const \; . \qquad (6)$$

The coefficients $d_\alpha(\gamma)$ can be evaluated analytically. In the Ising case the following symmetry holds: $d_x(1) = -c_x(1)$, $d_y(1) = -c_y(1)$. In the whole interval $0 < \gamma < 1$ the accelerations Λ_x and Λ_y exhibit a very similar scaling behavior in the proximity of the quantum critical point, obeying in all cases Eq. (6). The coefficients $d_{x,y}(\gamma)$ in Eq. (6) are related to the coefficients $c_{x,y}(\gamma)$ in Eq. (3) by a simple translation of minus the latter by an amount that decreases monotonically with the value of the anisotropy and vanishes when the latter reaches its maximum $\gamma = 1$. As an example, at $\gamma = 0.5$ the coefficients $d_x(0.5)$ and $d_y(0.5)$ are $d_x(0.5) = -c_x(0.5) + 0.239$ and $d_y(0.5) = -c_y(0.5) + 0.239$.

The reason why $d\Lambda_z/d\lambda$ does not exhibit scaling in the Ising model can be understood by comparing with its behavior at different values of the anisotropy. One finds that $d\Lambda_z/d\lambda$ scales logarithmically, just like $d\Lambda_x/d\lambda$ and $d\Lambda_y/d\lambda$, when approaching the quantum critical point in the whole interval $0 < \gamma < 1$. At first sight, this difference in the behavior of the acceleration Λ_z for the Ising model versus the other anisotropic cases would seem to imply a violation of the universality principle. However, studying the behavior of Λ_z as a function of λ when the anisotropy parameter γ approaches the Ising limit, one finds that $d\Lambda_z/d\lambda$ still obeys Eq. (6), but with a coefficient $d_z(\gamma)$ that decreases monotonically and vanishes



FIG. 4: Behavior of the first derivatives of the velocities $\Lambda_x = \Lambda_y$ and of the velocity Λ_z as functions of λ when approaching the critical point λ_c for the isotropic XY model ($\gamma = 0$). Red line: $d\Lambda_z/d\lambda$; black line: $d\Lambda_x/d\lambda = d\Lambda_y/d\lambda$.

exactly at $\gamma = 1$.

In the case of the isotropic XY model, we enter in a different class of universality and the behavior of the derivatives of the diffusion coefficients changes drastically compared to the anisotropic cases. Looking at Eq. (5) and taking into account that rotational symmetry implies $G_{xzx} = G_{yzy}$, we obtain $\Lambda_x = \Lambda_y = -G_z - G_{xzx}/2$. At variance with the behavior of the derivatives of the three energy gaps (compare Fig.2 and Fig.4), *all* the derivatives of the three accelerations $\Lambda_x = \Lambda_y$ and Λ_z diverge when approaching the critical point with a behavior analogous to that of the derivative of the energy gap ΔE_z (see Eq. (4)):

$$\frac{d\Lambda_\alpha}{d\lambda} = d'_\alpha(\lambda^2 - \lambda_c^2)^{-1/2} \, , \tag{7}$$

with $d'_x = d'_y = -1/\sqrt{2\pi^2}$ and $d'_z = -\sqrt{2}/\pi$.

In conclusion, we have introduced and studied observable quantities, either static or dynamic, associated with single qubit unitary operations, that are able to characterize quantum phase transitions in spin systems. This provides an alternative, quantum-informatic point of view complementary to the traditional one in terms of correlation functions. Moreover, at variance with the characterization of quantum critical points based on measures of entanglement, the quantities that we have introduced are, in principle, amenable to *direct* experimental observation. It is matter for future studies to establish whether this formalism can be useful for the characterization of other quantum critical systems and models, and in different research contexts. In this last respect, it would be particularly interesting to investigate possible connections with quantum information transfer, information speed, and quantum speed limits in critical systems [16, 17, 18].

We thank L. Amico, F. de Pasquale, and R. Fazio for useful discussions. Financial support from MIUR, Coherentia CNR-INFM, and INFN is acknwoledged.

---

[1] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, 2000).

[2] M. B. Hastings, Phys. Rev. Lett. **93**, 126402 (2004); *ibidem*, Phys. Rev. Lett. **93**, 140402 (2004).

[3] A. Osterloh, L. Amico, G. Falci, and R. Fazio, Nature **416**, 608 (2002).

[4] T. J. Osborne and M. A. Nielsen, Phys. Rev. A **66**, 032110 (2002).

[5] P. Calabrese and J. Cardy, J. Stat. Mech. (2004) P06002.

[6] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).

[7] T. Roscilde, P. Verrucchi, A. Fubini, S. Haas, and V. Tognetti, Phys. Rev. Lett. **93**, 167203 (2004); *ibidem*, Phys. Rev. Lett. **94**, 147208 (2005).

[8] S.-J. Gu, H.-Q Lin, and Y.-Q. Li, Phys. Rev. A **68**, 042330 (2003).

[9] Y. Chen, P. Zanardi, Z. D. Wang, and F. C. Zhang, quant-ph/0407228.

[10] S.-J. Gu, G.-S. Tian, and H.-Q. Lin, quant-ph/0511243.

[11] L.-A. Wu, M. S. Sarandy, D. A. Lidar, and L. J. Sham, quant-ph/0512031.

[12] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).

[13] E. Lieb, T. Schultz, and D. Mattis, Ann. Phys. (N.Y.) **16**, 407 (1961).

[14] E. Barouch, B. McCoy, and M. Dresden, Phys. Rev. A **2**, 1075 (1970); E. Barouch and B. McCoy, Phys. Rev. A **3**, 786 (1971).

[15] P. Pfeuty, Ann. Phys. (N.Y.) **57**, 79 (1970).

[16] S. Bose, Phys. Rev. Lett. **91**, 207901 (2003).

[17] V. Giovannetti, S. Lloyd, and L. Maccone, Phys. Rev. A **67**, 052109 (2003).

[18] M. J. Hartmann, M. E. Reuter, and M. B. Plenio, quant-ph/0511185.